

GRAS Notice (GRN) No. 520

<http://www.fda.gov/Food/IngredientsPackagingLabeling/GRAS/NoticeInventory/default.htm>

GR



ORIGINAL SUBMISSION

000001

April 28, 2014

Office of Food Additive Safety  
Division of Biotechnology and GRAS Notice Review, (HFS-255)  
Center for Food safety and Applied Nutrition  
Food and Drug Administration  
5100 Paint Branch Parkway  
College Park, MD 20740

RECEIVED  
MAY 14 2014  
OFFICE OF  
FOOD ADDITIVE SAFETY  
GRN 000520

To whom it may concern,

In accordance with the 21 CFR 170.36 (62 FR18937; April 17, 1997), MitoGene Beverage Company, LLC hereby provides notice of a claim that the use of **molecular hydrogen** (H<sub>2</sub>) solubilized in water is generally recognized as safe (GRAS) and therefore exempt from the premarket approval requirement of the Federal Food, Drug and Cosmetic Act. Based upon scientific data and information, including that in the attached documents, MitoGene Beverage Company has determined, that molecular hydrogen solubilized in water is general recognized as safe. The data and information that are the basis for this GRAS determination are available for the Food and Drug Administration's (FDA) review and copying at reasonable times at the address specified below in the "Information About the Notifier" section, or will be sent to the FDA upon request. As required, three copies of the notification, including supporting documentation, are provided.

If you have any questions regarding this notification, please feel free to contact any one of the submitters listed here. Thank you very much for your time and consideration.

Sincerely,

(b) (6)

04/28/2014

Dr. Cody C. Cook, Ph.D., M.D.  
Founder, MitoGene, LLC  
Cody.Cook@Gmail.com  
501-258-3079

(b) (6)

Apr. 28, 2014

Dr. Drew R. Jones, Ph.D.  
Co-Founder, MitoGene, LLC  
DrewRJones@Gmail.com  
505-250-9921

Enclosures

000002

### **Information About The Notifier**

MitoGene Beverage Company, LLC  
Cody Cook (Manager)  
Drew Jones (Manager)

401 S. Cedar St  
Little Rock, Arkansas 72205  
OR:  
1544 Avilla Vincintage Road  
Alexander, Arkansas 72002

### **Intended Use**

This GRAS notification serves to delineate the intended use of molecular hydrogen up to a concentration of 2.14% (vol H<sub>2</sub> (gas)/vol H<sub>2</sub>O (liq)) (saturation) in pure drinking water, flavored beverages, and soda drinks for the purpose of preventing oxidation to the beverages, beverage containers, and the consumer.

### **Identity of Notified Substance**

#### *Hydrogen Gas*

Synonyms for hydrogen include dihydrogen, ortho-hydrogen/para-hydrogen, and most commonly, molecular hydrogen. Hydrogen has an empirical formula of H, and a structural formula of H<sub>2</sub>. The CAS number of hydrogen is 1333-74-0.

Molecular weight: 2.016 g/mol

Specific volume (1.013 bar and 21 °C, 70 °F): 11.986 m<sup>3</sup>/kg

Solubility in water (1.013 bar and 0 °C, 32 °F): 0.0214 mL H<sub>2</sub> (gas)/mL H<sub>2</sub>O (liq)

Concentration in air: 0.00005% (vol)

*Description of Hydrogen Gas:* Hydrogen is a non-toxic, flammable, lighter than air gas, which reacts with oxidizing compounds. In terms of environmental and human health, the main risk of hydrogen exposure is due to its flammability; the gas is flammable in air between 4-75% by volume. The set exposure limits to hydrogen are 100% by volume due to oxygen depletion by simple asphyxiation. As a cryogenic fluid, hydrogen exposure can cause frostbite.

According to manufacturer material safety data sheets (MSDS), and a wealth of scientific literature, hydrogen has no known acute or chronic toxicity, carcinogenic effects, mutagenic effects, or reproductive toxicity. When dissolved in water, hydrogen has a rating of 0/0/0 for health/flammability/instability on the Hazardous Material Information System (U.S.A) indicating that it represents no significant risk toward human health, is not flammable, and is chemically stable.

000003

### **Method of manufacture**

*Manufacturing of Hydrogen Gas:* Food grade molecular hydrogen gas, according to the intended application, is produced industrially through steam reformation of natural gas (NexAir). Trace contaminants include oxygen (O<sub>2</sub>), water vapor (H<sub>2</sub>O), total hydrocarbon (THC), argon (Ar), carbon monoxide (CO), carbon dioxide (CO<sub>2</sub>), and nitrogen (N<sub>2</sub>). None of the trace contaminants are toxic at the specified levels or in the intended use.

### **Specifications – Good Manufacturing Practices**

Food grade hydrogen should be High Purity (99.995%) or greater. To establish acceptable contaminant levels in food grade hydrogen for good manufacturing practices, the company has assessed the currently accepted limits of similarly used food grade gases, e.g., carbon dioxide and nitrogen. Based on these data (Figures 3, 4), we have established levels for food grade hydrogen (Figure 2). All executions of the described intended use in this notification will follow these good manufacturing practices.

### **Applicable Conditions of Use**

The chemical properties of hydrogen make it ideally suited for several applications within the food and beverage industry. Molecular hydrogen is non-toxic and is a potent antioxidant, making it a valuable preservative. The current intended application is the use of solubilized H<sub>2</sub> in pure drinking water, flavored beverages, and soda drinks. The presence of H<sub>2</sub> in such beverages serves as an antioxidant and protecting the beverage and beverage delivery system (i.e. the can) from oxidation.

Hydrogen (H<sub>2</sub>)-enriched beverages (including drinking water, flavored water, and soft-drinks), are produced by introducing high purity (99.995%) hydrogen gas into purified drinking water under high-pressure (~ 100 psi), loosely analogous to the carbonation process for carbonated beverages. This process solubilizes hydrogen gas at the saturation limit of water. Due to the chemical solubility of hydrogen in water, the concentration in such beverages reaches up to 0.0214 mL H<sub>2</sub> (gas)/mL H<sub>2</sub>O (liq). In contrast, carbonated beverages may contain up to 1.72 mL CO<sub>2</sub> (gas)/mL H<sub>2</sub>O (liq). The resulting product contains hydrogen at a concentration well below the lower flammability limit. In this form, dissolved hydrogen is completely non-toxic, and non-flammable.

Because of hydrogen's biological safety profile, hydrogen-enriched beverages are suitable for persons of all ages and medical conditions under the described application conditions (consumption of up to 2 L/day of 0.0214 mL H<sub>2</sub> gas/mL H<sub>2</sub>O). As of this date, there is no evidence in the scientific literature that H<sub>2</sub>-enriched beverages should not be consumed by any segment of the population.

### **Self-limiting Levels of Use**

The solubilization of H<sub>2</sub> in a beverage limits the access of the molecule to oxygen such that it is non-flammable in a liquid medium. Due to the chemical solubility of hydrogen in water (Figure 1), hydrogen may reach a maximum concentration of 0.0214 mL H<sub>2</sub> gas/mL H<sub>2</sub>O (liq), corresponding to 1.8 ppm. This concentration of H<sub>2</sub> in water is well below the lower flammability limit of molecular hydrogen (4% H<sub>2</sub> (gas)) in air and can't be concentrated further. Upon exposure to the air, the dissolved hydrogen slowly diffuses out of solution (with a half-life of ~2 hours) and

into the air, becoming more dilute. Notably, hydrogen is naturally present in air at 0.000055% (v/v).

### **Dietary Exposure**

Assuming a maximum consumption of 2 L hydrogen enriched beverages per day, total hydrogen exposure is equal to 3.2 mg of hydrogen. Given that the only health risk of hydrogen exposure is due to suffocation, these consumption levels are generally regarded as safe by scientists, clinicians, and as verified through scientific proceedings.

### **History of Use: Hydrogen as an Antioxidant/Food Preservative**

The international community recognizes the value of hydrogen as a food preservative, antioxidant, and packaging leak indicator. Molecular hydrogen is used in the European Union, under the category of a packaging gas E949 ([http://www.food.gov.uk/policy-advice/additivesbranch/enumberlist#h\\_7](http://www.food.gov.uk/policy-advice/additivesbranch/enumberlist#h_7)). In this context, liquids and food stuffs are exposed to and stored in the presence of H<sub>2</sub> gas for long periods of time, whereby hydrogen diffuses through and permeates the product. Hydrogen becomes absorbed such that it protects the product and/or container from oxidative degradation. Hydrogen gas is ideal for these purposes because of its large specific volume, low molecular weight, and non-toxic profile.

### **Comprehensive Discussion of the Scientific Proceedings used for the GRAS Determination**

The determination that animal and human consumption of H<sub>2</sub> in beverages is GRAS is based on scientific data, information, methods, and principles that have been collected from the publically available scientific literature. Reports of the consumption of H<sub>2</sub> in animal and human studies are unanimous in the standing that H<sub>2</sub> is GRAS. The reports have been extensively published in primary, peer-reviewed scientific journals of the highest tier (Dole, Wilson et al. 1975; Ohsawa, Ishikawa et al. 2007), and supported by the secondary scientific literature such as scientific review articles (Wood and Gladwin 2007). The consumption of H<sub>2</sub> saturated water has been extensively studied (Shirahata, Kabayama et al. 1997; Kim and Kim 2006). Electrolyzed-reduced water is characterized with a highly negative redox potential, in addition to, high dissolved H<sub>2</sub> content. Consumption of hydrogen-enriched water has been extensively studied (Kajiyama, Hasegawa et al. 2008; Ohsawa, Nishimaki et al. 2008; Sato, Kajiyama et al. 2008; Fu, Ito et al. 2009; Fujita, Seike et al. 2009; Kajiya, Silva et al. 2009; Nagata, Nakashima-Kamimura et al. 2009; Nakashima-Kamimura, Mori et al. 2009; Cardinal, Zhan et al. 2010; Liu, Liu et al. 2011). The consumption of H<sub>2</sub> has been assessed across a variety of disease models and many research contexts including; oxidative stress markers, glucose metabolism, diabetes, inflammatory bowel disease, and pathogenic events (Kamimura, Nishimaki et al. 2011., Kajiyama, Hasegawa et al. 2008; Nakao, Toyoda et al. 2010., Kajiya, Silva et al. 2009; Nakashima-Kamimura, Mori et al. 2009).

### **Disposition of Molecular Hydrogen in Humans**

The pharmacokinetic properties of hydrogen-enriched beverages have been evaluated by the medical and scientific communities over decades of research in many fields of study. The

following data review and highlight key scientific studies demonstrating the scientific consensus that the intended application described in this document is generally regarded as safe.

### **Absorption**

After consuming H<sub>2</sub>-enriched beverages, H<sub>2</sub> is rapidly and completely absorbed in healthy adult volunteers that have fasted, as determined by hydrogen breath tests. The concentration of H<sub>2</sub> in exhaled breath reaches a peak (35-to-56.8 ± 27.8 ppm) at 15 minutes in a dose-dependent manner (Kajiya, Hasegawa et al. 2008; Shimouchi, Nose et al. 2009). H<sub>2</sub> reaches peak absorption within ~15 minutes following oral consumption (Cardinal, Zhan et al. 2010). It is generally reported that acute ingestion of H<sub>2</sub>-enriched water rapidly increases the H<sub>2</sub> partial pressure in the alimentary tract, and sub-mucosal alimentary vessels. Kajiya et al. and others have reported that the concentration of H<sub>2</sub> in the stomach of animal models, that were directly administered H<sub>2</sub>-enriched water by catheter, increased within 5 minutes (503.8 ± 160.3 μM, 1.0 ppm) until the concentration began to decrease by 10 minutes (200.8 ± 106.6 μM, 0.40 ppm) and return to baseline levels within one hour (Kajiya, Silva et al. 2009). The same report noted that the concentration of H<sub>2</sub> in the colon gradually increased to its highest level (88.0 ± 26.5 μM, 0.18 ppm) at 30 minutes, presumably by a combination of cross-epithelial diffusion (Lemke, van Alen et al. 2001) and the vascular-based transport processes (Bond, Levitt et al. 1975), until 50 minutes, when the concentration of H<sub>2</sub> returned to the exogenous baseline levels already present in the colon (58.0 ± 15.1 μM, 0.12 ppm) (Kajiya, Silva et al. 2009). Nagata et al. has reported that the incorporation of (0.8 mM, 1.6 ppm) hydrogen-enriched water directly into the stomach of a rat (3.5 mL per 230 g, 15 mL/kg) increased the H<sub>2</sub> concentration in blood within 3 minutes (Nagata, Nakashima-Kamimura et al. 2009), while a similar study has reported that the concentration of H<sub>2</sub> in the blood is increased by 3.7-fold and 7.6-fold in the fed and fasted state (Nakashima-Kamimura, Mori et al. 2009). It should also be noted that hydrogen gas has been administered to animals and humans through intraperitoneal injection, intravenous injection, and via eye drops with similar results to oral consumption (Mao, Zheng et al. 2009; Nakayama, Kabayama et al. 2009; Zheng, Mao et al. 2009; Oharazawa, Igarashi et al. 2010; Zheng, Liu et al. 2010; Chen, Sun et al. 2011; Ji, Hui et al. 2011).

### **Distribution**

H<sub>2</sub> is electronically neutral and is much smaller than molecular oxygen (O<sub>2</sub>). These properties allow H<sub>2</sub> to penetrate the cellular and intracellular membranes of cells and organelles explaining its large volume of distribution. The highly diffusible and lipid soluble nature of H<sub>2</sub> enables it to reach equilibrium nearly instantly with respect to blood and tissue. This fast diffusion results in H<sub>2</sub> being distributed to tissues in proportion to the regional blood flow of the tissue. These properties have been exploited in the clinic, making H<sub>2</sub> a powerful tool for measuring arterial and venous blood flow (Aukland, Bower et al. 1964). When consumed orally via enriched beverages, H<sub>2</sub> is distributed throughout the total volume of water in the body. Currently there are still technical challenges in detecting low levels of H<sub>2</sub> within tissues (Fujita, Seike et al. 2009). During the distribution phase, H<sub>2</sub> is diluted by 160 fold between the stomach and blood (Nagata, Nakashima-Kamimura et al. 2009). There is an even greater expected dilution factor between the blood and tissue, suggesting that the concentration of H<sub>2</sub> in distal tissues is below the detection limit of even the most sensitive analytical methods currently available. In chronic

exposure studies (60 days), the level of H<sub>2</sub> in the circulation of animals is comparable to that of naive animals (Cardinal, Zhan et al. 2010). These results confirm that H<sub>2</sub> does not accumulate in the body after consumption of H<sub>2</sub>-enriched beverages. Further, the pharmacokinetic parameters did not change during the course of the study.

### **Metabolism**

The metabolism of H<sub>2</sub> is important from both anabolic and catabolic perspectives. H<sub>2</sub> is a natural component of abdominal gas (Hammer 1993) where it is reportedly catabolized by colonic microflora into still unconfirmed metabolites such as methane (Christl, Murgatroyd et al. 1992). Endogenous H<sub>2</sub> is also generated through anabolic pathways carried out exclusively by numerous strains of colonic microflora (Reth 2002). Fermentation is also increased in response to a diet of numerous foods with un-absorbable carbohydrates, such as milk, commercial dietary fiber, soybean flour, and oligofructose in a number of vegetables, fruits, and whole grains (O'Brien, Thompson et al. 1987; Kondo and Nakae 1996; Suarez, Springfield et al. 1999; Cherbut 2002; Shimouchi, Nose et al. 2009). For example, the ingestion of milk exhibits a delayed and sustained increase of breath hydrogen (164 ppm) in subjects with hypolactasia for up to 540 minutes, in comparison to the relative (60 min) transient increase in hydrogen (2-to-55 ppm) that is noted with H<sub>2</sub>-enriched water (Kajiyama, Hasegawa et al. 2008; Nagata, Nakashima-Kamimura et al. 2009; Shimouchi, Nose et al. 2009).

### **Excretion/Elimination**

Following absorption via oral consumption, the concentration of H<sub>2</sub> in the blood decreases to baseline levels (11.2 ± 6.5 ppm) within 2.5 hours (Kajiyama, Hasegawa et al. 2008). The major route of H<sub>2</sub> elimination is through the respiratory system due to the low rate of metabolism, and high volatility of the compound (Shimouchi, Nose et al. 2009). The vapor-liquid equilibrium constant for H<sub>2</sub> drives hydrogen into the vapor phase, facilitating gas exchange of H<sub>2</sub> from the blood to the alveolar ducts. Once in the gas phase, hydrogen is uni-directionally expelled from the body upon exhalation due to the ~40,000-fold difference in concentration of H<sub>2</sub> between the blood and the atmosphere.

**(III). Safety Profile of the Consumption of H<sub>2</sub>:** The safety profile of the consumption of H<sub>2</sub> is a reflection of the well-defined characteristics of the properties of the gas with respect to the human body. The role of H<sub>2</sub> in the body has been studied extensively (Ohsawa, Ishikawa et al. 2007; Fujita, Seike et al. 2009)(Sato, Kajiyama et al. 2008)(Bjelakovic, Nikolova et al. 2007; Ohsawa, Ishikawa et al. 2007; Wood and Gladwin 2007). Throughout the myriad animal models and human studies which have exploited hydrogen consumption, either as a non-flammable gas (<4%) or in H<sub>2</sub>-enriched water (< 0.8 mM, 1.6 ppm), there are no reports to-date of any toxicity or deleterious changes in physiological parameters (e.g., body temperature, heart rate) associated with administration of H<sub>2</sub>. (Ohsawa, Ishikawa et al. 2007; Hayashida, Sano et al. 2008; Kajiyama, Hasegawa et al. 2008; Ji, Liu et al. 2010; Nakao, Toyoda et al. 2010; Xie, Yu et al. 2010; Xie, Yu et al. 2010; Huang, Xie et al. 2011). It is important to note that the short-term and long-term consumption of H<sub>2</sub> in animal and human studies has been evaluated in animal models and human studies that are subjected to a wide variety of (patho)physiologies of organ

damage, and multiple-organ damage. The lack of negative interactions with the consumption of H<sub>2</sub> in these models of poor health further demonstrates that H<sub>2</sub> is Generally Regarded As Safe.

The safety profile for the human consumption of H<sub>2</sub> has been further supported with 17 year-long studies on cells, mice, monkeys, and deep sea divers (COMEX HYDRA program, Marseille). Extremely high concentrations of H<sub>2</sub> (49%) have been utilized in hydroliox for very deep technical diving ventures to shorten decompression time, prevent decompression sickness, avert nitrogen narcosis, and reduce airway pressure (Abraini, Gardette-Chauffour et al. 1994). The hydroliox mixture consisted of hydrogen, helium and oxygen at relative concentrations of 49% H<sub>2</sub>, 50% He<sub>2(gas)</sub> and 1% O<sub>2(gas)</sub>, respectively. The absorption of H<sub>2</sub> in these extremely high concentrations has not been associated with toxic effects, including no reports of intestinal motility changes (Lafay, Barthelemy et al. 1995; Fontanari, Badier et al. 2000). The high concentration of H<sub>2</sub> that has been utilized in inhalation and diver studies suggest that 16 mM in blood could be safe (Fontanari, Badier et al. 2000), confirming that the concentrations reached with the consumption of H<sub>2</sub> in beverages is GRAS.

Moreover, endogenous breath hydrogen levels fluctuate markedly during daily life, ranging from several ppm to 50 ppm or more (Sone, Tanida et al. 2000). The concentration of H<sub>2</sub> in the circulation of humans is enhanced by exercise, physical stress, and mental stress (Kondo and Nakae 1996; Suarez, Springfield et al. 1999). Thus, the hydrogen-producing capability of H<sub>2</sub>-enriched water is within the range of H<sub>2</sub> that is produced when people with hypolactasia consume milk.

**Scientific Conclusion that Hydrogen-Enriched Beverages are GRAS:** The collective review of the scientific literature shows a unanimous acknowledgment that H<sub>2</sub>, one of the most well understood molecules, is a generally recognized as safe alternative to the commercialized antioxidants in beverages based on the premise that the gas is easily available in pure form and without toxicity. Overall, no toxicity of any kind has been attributed to the *ad libitum* consumption of H<sub>2</sub> in the diet. Similarly, no contradictions have been noted with any other chemically or pharmacologically related or non-related substance in the diet of animals or humans. The scientific literature is devoid of any report of investigations or other information that may appear to be inconsistent with the MitoGene, LLC self-determination that above described intended use of hydrogen is GRAS.

**Signatures**

(b) (6)



04/28/2014

Cody C. Cook, Ph.D., M.D.  
Manager, MitoGene Beverage Company, LLC


(b) (6)



April 28, 2014

Drew R. Jones, Ph.D.  
Manager, MitoGene Beverage Company, LLC

## Figures



**Figure 1: Solubility of Hydrogen Gas in Degassed, Distilled Water.** Data adapted from (Thomas E. Crozier 1974).



Gas Technologies  
Medical Technologies  
Safety Technologies  
Welding Technologies

January 2, 2013

Dear Customer:

This is your Certificate of Analysis:

Product: Hydrogen

Sample Serial Number: (b) (6)

Lot Number: (b) (6)

Component	Min Purity	Analytical Results	Analytical Method
Oxygen	<1.0 ppm	<1.0 ppm	Electrochemical Cell (Delta F)
Moisture	<3.0 ppm	<2.0 ppm	Meeco
Total Hydrocarbons	<0.5 ppm	<0.5 ppm	THC Analyzer
Nitrogen	<6.0 ppm	< 6.0 ppm	GC-TCD
Assay		> 99.999%	

(b) (6)

Jason Spires  
Lab Analyst

nexAir, LLC · 1259 N McLean Blvd · Memphis, TN 38108 · (901) 729-5540 · (800) 294-8062 · (901) 729-5575 fax · www.nexair.com

Figure 2: Certificate of Analysis for Food Grade Hydrogen

000011



November 19, 2013

nexAir  
PO Box 161182  
Memphis, TN 38116

Praxair # 8512262

Dear Customer,

This is to advise you that liquid Carbon Dioxide supplied by Praxair, Inc. to your facility at 1211 North McLean Blvd, Memphis, TN 38108 meets or exceeds the Food Grade specifications as follows:

Minimum Assay: 99.8%	Total Sulfur: 0.5 ppm
Acetaldehyde: 0.5 ppm	Total Hydrocarbon content (as methane) : 50 ppm
Carbon Monoxide: 10 ppm	Hydrogen Sulfide: 0.5 ppm
Carbonyl Sulfide: 0.5 ppm	Non-Volatile Residue: 10 ppm
Nitric Oxide + Nitrogen Dioxide: 5 ppm	Odor/Taste: Free of foreign odor or taste
Oxygen: 50 ppm	Moisture: 20 ppm
Sulfur Dioxide: 5 ppm	

The CGA lists the typical use for this grade of product as food processing. Carbon Dioxide is listed in the Code of Federal Regulations 21 CFR 184.1240 as Generally Recognized as Safe (GRAS) as a direct human food ingredient (see attached).

Sincerely,

Tim Lowry  
Praxair, Inc  
Senior Account Manager

**Figure 3: Reference Ranges of Contaminants for Food Grade Carbon Dioxide**

**000012**



April 11, 2013

nexAir, LLC  
PO Box 161182  
Memphis, TN 38186

Praxair # 8507527

Dear Customer:

This is to advise you that the liquid nitrogen supplied by Praxair, Inc. delivered to nexAir at 1211 N McLean Blvd, Memphis, TN 38108 has a purity guarantee of 99.998% by difference and meets or exceeds the following specifications:

<u>COMPONENT</u>	<u>SPECIFICATION</u>
Oxygen	8.0 PPM
Moisture	4.0 PPM
Carbon Monoxide	10.0 PPM
Odor	None detected

Nitrogen is listed in the Code of Federal Regulations 21 CFR 184.1540 as Generally Recognized as Safe (GRAS) as a direct human food ingredient. See attached. Also, the Compressed Gas Association lists typical uses for this grade of product as general industrial usage, process gases, inerting, food, and heat treatment applications.

Sincerely,

(b) (6)

Tim Lowry  
Praxair, Inc.  
NAIG Senior Account Manager

Page 1 of 2

Figure 4: Reference Ranges of Contaminants for Food Grade Nitrogen

000013

## Bibliography

Abraini, J. H., M. C. Gardette-Chauffour, et al. (1994). "Psychophysiological reactions in humans during an open sea dive to 500 m with a hydrogen-helium-oxygen mixture." J Appl Physiol **76**(3): 1113-1118.

Aukland, K., B. F. Bower, et al. (1964). "Measurement of Local Blood Flow with Hydrogen Gas." Circ Res **14**: 164-187.

Bjelakovic, G., D. Nikolova, et al. (2007). "Mortality in randomized trials of antioxidant supplements for primary and secondary prevention: systematic review and meta-analysis." JAMA **297**(8): 842-857.

Bond, J. H., Jr., M. D. Levitt, et al. (1975). "Investigation of small bowel transit time in man utilizing pulmonary hydrogen (H<sub>2</sub>) measurements." J Lab Clin Med **85**(4): 546-555.

Buchholz, B. M., D. J. Kaczorowski, et al. (2008). "Hydrogen inhalation ameliorates oxidative stress in transplantation induced intestinal graft injury." Am J Transplant **8**(10): 2015-2024.

Cai, J., Z. Kang, et al. (2008). "Hydrogen therapy reduces apoptosis in neonatal hypoxia-ischemia rat model." Neurosci Lett **441**(2): 167-172.

Cardinal, J. S., J. Zhan, et al. (2010). "Oral hydrogen water prevents chronic allograft nephropathy in rats." Kidney Int **77**(2): 101-109.

Chen, H., Y. P. Sun, et al. (2011). "The effects of hydrogen-rich saline on the contractile and structural changes of intestine induced by ischemia-reperfusion in rats." J Surg Res **167**(2): 316-322.

Cherbut, C. (2002). "Inulin and oligofructose in the dietary fibre concept." Br J Nutr **87 Suppl 2**: S159-162.

Christl, S. U., P. R. Murgatroyd, et al. (1992). "Production, metabolism, and excretion of hydrogen in the large intestine." Gastroenterology **102**(4 Pt 1): 1269-1277.

Di Camillo, M., V. Marinaro, et al. (2006). "Hydrogen breath test for diagnosis of lactose malabsorption: the importance of timing and the number of breath samples." Can J Gastroenterol **20**(4): 265-268.

Dole, M., F. R. Wilson, et al. (1975). "Hyperbaric hydrogen therapy: a possible treatment for cancer." Science **190**(4210): 152-154.

Fontanari, P., M. Badier, et al. (2000). "Changes in maximal performance of inspiratory and skeletal muscles during and after the 7.1-MPa Hydra 10 record human dive." Eur J Appl Physiol **81**(4): 325-328.

Fu, Y., M. Ito, et al. (2009). "Molecular hydrogen is protective against 6-hydroxydopamine-induced nigrostriatal degeneration in a rat model of Parkinson's disease." Neurosci Lett **453**(2): 81-85.

Fujita, K., T. Seike, et al. (2009). "Hydrogen in drinking water reduces dopaminergic neuronal loss in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson's disease." PLoS One **4**(9): e7247.

Fukuda, K., S. Asoh, et al. (2007). "Inhalation of hydrogen gas suppresses hepatic injury caused by ischemia/reperfusion through reducing oxidative stress." Biochem Biophys Res Commun **361**(3): 670-674.

Gharib, B., S. Hanna, et al. (2001). "Anti-inflammatory properties of molecular hydrogen: investigation on parasite-induced liver inflammation." C R Acad Sci III **324**(8): 719-724.

Hammer, H. F. (1993). "Colonic hydrogen absorption: quantification of its effect on hydrogen accumulation caused by bacterial fermentation of carbohydrates." Gut **34**(6): 818-822.

Hayashi, T., T. Yoshioka, et al. (2011). "Inhalation of hydrogen gas attenuates left ventricular remodeling induced by intermittent hypoxia in mice." Am J Physiol Heart Circ Physiol **301**(3): H1062-1069.

Hayashida, K., M. Sano, et al. (2008). "Inhalation of hydrogen gas reduces infarct size in the rat model of myocardial ischemia-reperfusion injury." Biochem Biophys Res Commun **373**(1): 30-35.

Huang, C. S., T. Kawamura, et al. (2011). "Hydrogen inhalation reduced epithelial apoptosis in ventilator-induced lung injury via a mechanism involving nuclear factor-kappa B activation." Biochem Biophys Res Commun **408**(2): 253-258.

Huang, Y., K. Xie, et al. (2011). "Beneficial effects of hydrogen gas against spinal cord ischemia-reperfusion injury in rabbits." Brain Res **1378**: 125-136.

James, A. M., H. M. Cocheme, et al. (2005). "Mitochondria-targeted redox probes as tools in the study of oxidative damage and ageing." Mech Ageing Dev **126**(9): 982-986.

Ji, Q., K. Hui, et al. (2011). "The effect of hydrogen-rich saline on the brain of rats with transient ischemia." J Surg Res **168**(1): e95-101.

Ji, X., W. Liu, et al. (2010). "Beneficial effects of hydrogen gas in a rat model of traumatic brain injury via reducing oxidative stress." Brain Res **1354**: 196-205.

Kajiya, M., M. J. Silva, et al. (2009). "Hydrogen mediates suppression of colon inflammation induced by dextran sodium sulfate." Biochem Biophys Res Commun **386**(1): 11-15.

Kajiyama, S., G. Hasegawa, et al. (2008). "Supplementation of hydrogen-rich water improves lipid and glucose metabolism in patients with type 2 diabetes or impaired glucose tolerance." Nutr Res **28**(3): 137-143.

Kamimura, N., K. Nishimaki, et al. (2011). "Molecular hydrogen improves obesity and diabetes by inducing hepatic FGF21 and stimulating energy metabolism in db/db mice." Obesity (Silver Spring) **19**(7): 1396-1403.

Kawamura, T., C. S. Huang, et al. (2010). "Inhaled hydrogen gas therapy for prevention of lung transplant-induced ischemia/reperfusion injury in rats." Transplantation **90**(12): 1344-1351.

Kim, M. J. and H. K. Kim (2006). "Anti-diabetic effects of electrolyzed reduced water in streptozotocin-induced and genetic diabetic mice." Life Sci **79**(24): 2288-2292.

Kondo, T. and Y. Nakae (1996). "Breath hydrogen and methane excretion produced by commercial beverages containing dietary fiber." J Gastroenterol **31**(5): 654-658.

Lafay, V., P. Barthelemy, et al. (1995). "ECG changes during the experimental human dive HYDRA 10 (71 atm/7,200 kPa)." Undersea Hyperb Med **22**(1): 51-60.

Lembcke, B., M. Honig, et al. (1980). "Different actions of neomycin and metronidazole on breath hydrogen (H<sub>2</sub>) exhalation." Z Gastroenterol **18**(3): 155-160.

Lemke, T., T. van Alen, et al. (2001). "Cross-epithelial hydrogen transfer from the midgut compartment drives methanogenesis in the hindgut of cockroaches." Appl Environ Microbiol **67**(10): 4657-4661.

Liu, S., K. Liu, et al. (2011). "Consumption of hydrogen water reduces paraquat-induced acute lung injury in rats." J Biomed Biotechnol **2011**: 305086.

Mao, Y. F., X. F. Zheng, et al. (2009). "Hydrogen-rich saline reduces lung injury induced by intestinal ischemia/reperfusion in rats." Biochem Biophys Res Commun **381**(4): 602-605.

Nagata, K., N. Nakashima-Kamimura, et al. (2009). "Consumption of molecular hydrogen prevents the stress-induced impairments in hippocampus-dependent learning tasks during chronic physical restraint in mice." Neuropsychopharmacology **34**(2): 501-508.

Nakao, A., D. J. Kaczorowski, et al. (2010). "Amelioration of rat cardiac cold ischemia/reperfusion injury with inhaled hydrogen or carbon monoxide, or both." J Heart Lung Transplant **29**(5): 544-553.

Nakao, A., Y. Toyoda, et al. (2010). "Effectiveness of hydrogen rich water on antioxidant status of subjects with potential metabolic syndrome-an open label pilot study." J Clin Biochem Nutr **46**(2): 140-149.

Nakashima-Kamimura, N., T. Mori, et al. (2009). "Molecular hydrogen alleviates nephrotoxicity induced by an anti-cancer drug cisplatin without compromising anti-tumor activity in mice." Cancer Chemother Pharmacol **64**(4): 753-761.

Nakayama, M., S. Kabayama, et al. (2009). "Biological effects of electrolyzed water in hemodialysis." Nephron Clin Pract **112**(1): c9-15.

O'Brien, J. D., D. G. Thompson, et al. (1987). "Action of centrally mediated autonomic stimulation on human upper gastrointestinal transit: a comparative study of two stimuli." Gut **28**(8): 960-969.

Oharazawa, H., T. Igarashi, et al. (2010). "Protection of the retina by rapid diffusion of hydrogen: administration of hydrogen-loaded eye drops in retinal ischemia-reperfusion injury." Invest Ophthalmol Vis Sci **51**(1): 487-492.

Ohsawa, I., M. Ishikawa, et al. (2007). "Hydrogen acts as a therapeutic antioxidant by selectively reducing cytotoxic oxygen radicals." Nat Med **13**(6): 688-694.

Ohsawa, I., K. Nishimaki, et al. (2008). "Consumption of hydrogen water prevents atherosclerosis in apolipoprotein E knockout mice." Biochem Biophys Res Commun **377**(4): 1195-1198.

Reth, M. (2002). "Hydrogen peroxide as second messenger in lymphocyte activation." Nat Immunol **3**(12): 1129-1134.

Sato, Y., S. Kajiyama, et al. (2008). "Hydrogen-rich pure water prevents superoxide formation in brain slices of vitamin C-depleted SMP30/GNL knockout mice." Biochem Biophys Res Commun **375**(3): 346-350.

Schneider, A. R., K. Jepp, et al. (2007). "The inulin hydrogen breath test accurately reflects oro-caecal transit time." Eur J Clin Invest **37**(10): 802-807.

Shimouchi, A., K. Nose, et al. (2009). "Breath hydrogen produced by ingestion of commercial hydrogen water and milk." Biomark Insights **4**: 27-32.

Shirahata, S., S. Kabayama, et al. (1997). "Electrolyzed-reduced water scavenges active oxygen species and protects DNA from oxidative damage." Biochem Biophys Res Commun **234**(1): 269-274.

Sone, Y., S. Tanida, et al. (2000). "Everyday breath hydrogen excretion profile in Japanese young female students." J Physiol Anthropol Appl Human Sci **19**(5): 229-237.

Suarez, F. L., J. Springfield, et al. (1999). "Gas production in human ingesting a soybean flour derived from beans naturally low in oligosaccharides." Am J Clin Nutr **69**(1): 135-139.

Swagerty, D. L., Jr., A. D. Walling, et al. (2002). "Lactose intolerance." Am Fam Physician **65**(9): 1845-1850.

Thomas E. Crozier, S. Y. (1974). "Solubility of hydrogen in water, sea water, and sodium chloride solutions." Journal of Chemical and Engineering Data **19**(3): 2.

Wood, K. C. and M. T. Gladwin (2007). "The hydrogen highway to reperfusion therapy." Nat Med **13**(6): 673-674.

Xie, K., Y. Yu, et al. (2010). "Protective effects of hydrogen gas on murine polymicrobial sepsis via reducing oxidative stress and HMGB1 release." Shock **34**(1): 90-97.

Xie, K., Y. Yu, et al. (2010). "Hydrogen gas improves survival rate and organ damage in zymosan-induced generalized inflammation model." Shock **34**(5): 495-501.

Zheng, J., K. Liu, et al. (2010). "Saturated hydrogen saline protects the lung against oxygen toxicity." Undersea Hyperb Med **37**(3): 185-192.

Zheng, X., Y. Mao, et al. (2009). "Hydrogen-rich saline protects against intestinal ischemia/reperfusion injury in rats." Free Radic Res **43**(5): 478-484.

**SUBMISSION END**

**000019**